



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

BRIEFER ARTICLES.

New localities.—The following are some localities for plants, not given in Gray's Manual, 6th edition:

Salsola kali tragus was first found and reported in Illinois, by myself, at Polo, Aug. 14th. Since then I have found it at Oregon, Savanna, and Chicago. Letters to the Experiment Station also give it as occurring at fourteen other places. At most places it was undoubtedly introduced by cattle trains from the north-west. So far it has been found only in the northern part of the state, and chiefly along the railroads. At Savanna, and possibly at some places in Chicago, this plant undoubtedly occurred last year, as two or three thousand plants were found.

Grindelia squarrosa is given as occurring at Evanston only, in this state. Several specimens of this were found by me at Polo, Oregon, and Savanna. These towns are on the Chicago, Burlington and Northern railroad, and the plants were undoubtedly introduced by that road from the north.

At Savanna a few specimens of *Solanum triflorum* were discovered. This is given in the Manual as occurring in central Kansas and westward. This was also introduced by the railroad.

A single specimen of what appears to be *Verbascum nigrum* was found on the Experiment Station grounds at Champaign. This probably came through seed from Europe, as the plant is not given in the Manual as yet occurring in the United States.—G. P. CLINTON, *Champaign, Ills.*

Two Wisconsin Fungi.—*Uromyces minimus*, n. sp.—Hypophyllous. Uredosori light brown, teleutosori black, oblong or linear, soon naked. Uredospores globose or oval, light brown, echinulate, 12–19 μ in diameter, usually 14–16. Teleutospores brown, smooth, spheroidal oval or oblong, 14–22 \times 12–19 μ , usually 17–20 \times 15–17 μ , apex rounded, conical or occasionally truncate, very strongly thickened, the apical thickening often constituting nearly half the length of the spore; pedicels moderately stout, tinted, once to twice the length of the spore. Colorless clavate paraphyses present.—On *Muhlenbergia sylvestica* Torr. & Gr. Kenosha co., Wisconsin.

In the same station an æcidium occurs on *Cacalia reniformis* Muhl. which differs from the other æcidia on Compositæ and which I have not seen elsewhere, but no experiments have been made to demonstrate genetic relationship.